

Medical Oncology

Drug Information Sheet

Palbociclib (Ibrance): a drug to treat your cancer.

What should I avoid while receiving *Palbociclib (Ibrance)*?

- Do NOT use Echinacea.
- Do not eat grapefruit or drink grapefruit juice while taking *Palbociclib (Ibrance)*.
- Do not take St. John's Wort with this drug.

Dose and Schedule:

- You will take this medicine by mouth.
- *Palbociclib (Ibrance)* should be taken at the same time every day (morning or evening)
- Take *Palbociclib (Ibrance)* with food.
- Your dose may vary, but the usual dose of *Palbociclib (Ibrance)* should be taken once daily for 21 days followed by 7 days off-treatment. The dose may be adjusted by your care provider based on your individual needs.

- Swallow the tablets or capsules whole with water. Do not crush, open, break, chew or dissolve the medicine in water.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- If you have any unused *Palbociclib (Ibrance)*, do not throw it in the trash and do not flush it down the sink or toilet. Talk to your doctor or pharmacist about proper disposal of *Palbociclib (Ibrance)*.
- *Palbociclib (Ibrance)* has many drug interactions. Tell your doctor all the prescription medicines, over-the-counter medicines, vitamins, and herbal products you are taking.
- Your doctor may change your dose, briefly stop or completely stop treatment with *Palbociclib (Ibrance)* if you have side effects.
- **DO NOT START A NEW CYCLE WITHOUT TALKING TO YOUR DOCTOR FIRST.**

- Capsules should not be opened. If capsules are accidentally opened or damaged, do not inhale contents or allow contents to come into contact with the skin or mucous membranes.
- If the inside contents are touched or they get in the eyes, wash hands or eyes right away.
- If the inside contents are accidentally inhaled, move to fresh air and call your doctor right away.

If you miss a dose of *Palbociclib (Ibrance)*:

- Do not take an extra dose or two doses at one time.
- Return to normal schedule the following day.
- If vomiting occurs after taking a dose, take the next dose at the scheduled time.
- Write down if you miss a dose and let your doctor know about any missed doses.

Medical Oncology

Drug Information Sheet

Storage and Handling:	
<ul style="list-style-type: none"> • You will get this drug from a special pharmacy through the mail. We will help you through the process. • The drug must be kept in a secure place out of reach of children and pets, so no one else can take this drug. Never share <i>Palbociclib (Ibrance)</i> with anyone else. • Store <i>Palbociclib (Ibrance)</i> at room temperature in a dry place and away from light. Do not store in a bathroom. • Keep in original bottle: do NOT place tablets in a pill box. 	
<ul style="list-style-type: none"> • Whenever possible, give <i>Palbociclib (Ibrance)</i> to yourself and follow the steps below. If a family member, friend or caregiver gives <i>Palbociclib (Ibrance)</i> to you, they also need to follow these steps <ul style="list-style-type: none"> ○ Wash hands with soap and water ○ Put on gloves to avoid touching the medicine (gloves are not needed if you give <i>Palbociclib (Ibrance)</i> to yourself). ○ Do not let women who are pregnant or who may get pregnant touch the tablets without wearing gloves. ○ Carefully remove <i>Palbociclib (Ibrance)</i> from its package to a small disposable cup. ○ Do not lay the medicine on the counter. ○ Take the medicine immediately by mouth with water. ○ Remove gloves and do not use them for anything else. ○ Throw gloves and disposable cup in household trash. ○ Wash hands with soap and water. ○ Do not touch or handle crushed or broken tablets. 	

You **MAY OR MAY NOT** experience the side effects listed below

Possible Side Effects:	What to do:
Increased risk for infection due to decreased blood counts	<ul style="list-style-type: none"> • Avoid contact with people who have infections, colds or flu. • Wash your hands often. • Report chills or a fever of 100.4° or higher to your doctor right away, day or night. • Report any signs or symptoms of infection, such as cough, burning with urination, skin redness or sores. • Call your doctor before taking any medicine for a fever or chills.
Lower than normal red blood cells Anemia	Call you doctor if you have: <ul style="list-style-type: none"> ○ shortness of breath with activity ○ fatigue ○ dizziness ○ fast heartbeat or a heartbeat that does not feel normal ○ headache
Hair loss	<ul style="list-style-type: none"> • Talk to your nurse about a wig or other options. • Ask about options to help decrease hair loss. • Hair will grow back 1-3 months after stopping chemotherapy.

Medical Oncology

Drug Information Sheet

Possible Side Effects:	What to do:
Fatigue or weakness	<ul style="list-style-type: none"> • You may be more tired than usual or have less energy. • Stay as active as possible, but know it is okay to rest as needed. • Try to do some activity every day. • Plan your activities, and do them at a time of day when you feel more energetic. • Avoid driving or doing other tasks that call for you to be alert.
Nausea/vomiting Decreased appetite	<ul style="list-style-type: none"> • Take nausea medicine as instructed. Let your doctor know if it is not working for you. • Eat smaller, frequent meals. • Try liquid nutrition supplements (Boost, Ensure). • A dietitian is available to talk with you.
Stomatitis Sores in your throat or mouth (stomatitis) may occur a few days or a few weeks after medicine is given.	<ul style="list-style-type: none"> • Let your nurse or doctor know if you have sores in your mouth. • Avoid spicy, hot or cold foods. • Avoid using tobacco products, alcohol, and mouthwashes that contain alcohol. • Mix 1/4 teaspoon of baking soda and 1/8 teaspoon of salt with 2 cups of water. Rinse your mouth and gargle the mixture at least four times daily, especially after you eat. Do not swallow this mixture, just rinse and spit. • Use a soft toothbrush. • Your doctor may prescribe a special mouthwash for you to use.
Liver problems	<ul style="list-style-type: none"> • Lab results will be monitored. • Call your doctor if you have: <ul style="list-style-type: none"> ○ dark urine (tea colored) ○ severe right sided stomach pain ○ light colored stools ○ bruising or if you start to bleed easily ○ yellowing of skin or eyes ○ severe nausea or throwing up ○ unexplained weight gain ○ your belly feels big or bloated

Medical Oncology

Drug Information Sheet

Possible Side Effects:	What to do:
<i>Palbociclib (Ibrance)</i> will be in urine and stool for 48 hours.	<ul style="list-style-type: none"> Caregivers will need to handle soiled linens and clothing wearing gloves or using a bag as a barrier. Keep your soiled items separate from others' dirty laundry and wash right away. Women who are pregnant should not handle your soiled clothing or linens. Wash any skin that body waste has touched with soap and water. You may use the same toilet that you normally use. If the toilet or toilet seat becomes dirty with urine, stool or vomit, clean the surfaces before other people use the toilet Wash hands with soap and water after using the toilet. It is safe to hug and kiss.
Avoid pregnancy	<ul style="list-style-type: none"> This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away. Use birth control that you can trust to prevent pregnancy while taking this drug and for 3 months after stopping this drug. If you or your sex partner gets pregnant while taking this drug or within 3 months after stopping this drug, call your doctor right away. May cause fertility problems.

Less common side effects that may happen

Less common side effects	What to do:
Inflammation of lungs	<ul style="list-style-type: none"> Call your doctor right away if you have: <ul style="list-style-type: none"> Severe shortness of breath or trouble breathing New cough or a cough that gets worse More production of mucus (sputum). Can be clear, white, yellowish-gray or green in color Wheezing Extreme tiredness Chest pain Fever of 100.4° or higher Chills Your doctor may place you on steroids.
These are not all of the side effects that may occur. If you have questions about side effects, call your doctor.	